



Testing the correct specification of a system of spatial dependence models for stock returns

Tim Kutzker¹ · Dominik Wied¹ 

Received: 7 June 2023 / Accepted: 10 October 2023
© The Author(s) 2023

Abstract

This paper provides two specification tests for the system of spatial autoregressive model of order m . We derive the theoretical limit distributions and show in a detailed Monte Carlo simulation study that the tests result in reasonable sized testing procedures with large power. In the empirical application, we analyze Euro Stoxx 50 returns in two different time spans, looking for insights how well models with different specifications of the spatial weighting matrices (local, country, industry and country-industry specific dependencies including interaction effects) fit to the data. The analyzes also demonstrate the ability of the tests to detect inaccurate Value-at-Risk forecasts.

Keywords Heteroscedasticity · Method of moments · Spatial dependence · Stock returns · Value-at-Risk

JEL Classification C12 · C51

1 Introduction

The purpose of this paper is a contribution to the literature of specification testing in spatial econometric models, focusing on systems of models. We consider a spatial autoregressive model in which a multivariate random variable is explained by spatially weighted lags of itself. Doing this, we allow for more than one spatial matrix and for cross-sectional heterogeneity of the error variances.

The authors acknowledge computational facilities by the Regional Computing Center of the University of Cologne via the DFG-funded High Performance Computing system CHEOPS (Grant INST 216/512/1FUGG). We are grateful to helpful comments from a referee.

✉ Tim Kutzker
tim.kutzker@gmail.com

Dominik Wied
dwied@uni-koeln.de

¹ Institute of Econometrics and Statistics, University of Cologne, Cologne, Germany

A classical reference for the question about the presence of global spatial dependence in a given data set is Moran's I (Moran 1950). This statistic is often used mainly in a descriptive way, but is also used for testing spatial dependence in linear and non-linear panel models (Kelejian and Prucha 2001). Li et al. (2007) provide an alternative measure to Moran's I. Other diagnostic tests for spatial dependence are the LM-tests by Anselin (1996) and the regression-based tests by Born and Breitung (2011). Su and Qu (2017) propose specification tests for SAR models.

However, neither of the mentioned tests consider the case of several time points. This is different for Baltagi et al. (2003) who propose LM-tests for spatial dependence and for Millo (2017) who proposes a randomization test in a factor-augmented panel. Kelejian and Piras (2016) propose a J-test procedure for testing a null model against non-nested alternatives for a fixed effects spatial panel data framework.

The model that we consider is slightly different from those considered in the approaches from the last paragraph. We focus on a repeated cross-sectional model, i.e. we have a SAR(m)-model for a fixed time point and stack these models for different time points. The model parameters can be estimated in a straightforward way by a two-step approach. This way, heterogeneity in the error variances can be incorporated easily.

Another related branch consists of testing procedures for selecting the best spatial weighting matrix from a set of potential candidates (Herrera et al. 2019; Kelejian and Piras 2011; LeSage and Pace 2014, e.g.). While our test can also be used for such model selection issues, the focus of specification testing is broader, since it tests the correctness of a model and its accompanying assumptions. We illustrate this in our application by first identifying spatial matrices modeling the spatial dependence of stock returns. We then use these spatial matrices to define different models, which we finally test with our model specification test.

It is precisely the connection between spatial dependence and stock returns that has received particular interest in the economics and finance literature in recent years. Ashgarian et al. (2013), for instance, investigate in which way stock market co-movements are determined by countries' economic and geographical relations. Tam (2014) analyzes equity market linkages in East Asia, Blasques et al. (2016) extend the spatial Durbin model by a time-varying spatial dependence parameter, Selan and Kalatzis (2017) analyze peer effects in Brazil.

A general observation is that the use of spatial and network statistical models is becoming more popular for modelling common stocks (both returns and volatility) rather than only stock markets of different countries. The role of geographical distance for modelling stock markets cross-correlation is quite debated, so the concept of "distance" is discussed and broadened using general definitions in the literature (cf. Fernández-Avilés et al. 2012; Fülle and Otto 2023; Mattera and Otto 2023; Zhou et al. 2020.)

More relevant for our work is Arnold et al. (2013) who consider a system of spatial autoregressive (SAR) models for stock returns in order to capture local dependencies and dependencies within industrial branches. Wied (2013) considers structural breaks in these models and Schmitt et al. (2016) combine the approach with local normalization techniques. Gong and Weng (2016) use the model for value at risk forecasts in the Chinese stock market. Catania and Billé (2017) generalize the SAR model with

autoregressive and heteroscedastic disturbances by including methods from score-driven models. Various empirical analyses in the aforementioned papers show that the SAR model is generally suitable for Value-at-Risk (VaR) forecasts and outperforms, e.g., the one-factor model.

The model in the present paper is a generalization of the model from Arnold et al. (2013) since we allow for an arbitrary fixed number of spatial matrices. We propose two methods on how to check the model fit, Note the fact that the model fits the data, implies that the spatial dependence structure incorporated in the model fits the data. Thus, we can interpret a specification test as a test for spatial dependence.

The basic idea stems from the model assumption that spatial weighting matrices capture all spatial dependence and that the remaining error terms are spatially uncorrelated. Therefore, we consider the model residuals to have the characteristic that if the covariance matrix of the residuals is basically diagonal, i.e. its off-diagonal elements are close to zero, the tests do not reject the null hypothesis of a correctly specified model. We derive the asymptotic distribution of our test statistics and show in simulations that the tests have reasonable power properties against sparse error term covariance matrices. In the simulations, we also account for conditional heteroscedasticity, a feature that is considered to be important if spatial models are used for VaR predictions (see, e.g., Zhang et al. 2018). In an empirical application on stock data from the Euro Stoxx 50, we test the model fit for different spatial weighting matrices and analyze in which sense the tests' results are related to the quality of VaR forecasts. We consider the time spans around the global financial crisis in 2008 and the COVID-19 crisis in 2020.

This paper is organized as follows: Sect. 2 describes the classical spatial autoregressive model, discusses its assumptions and efficient estimation procedures and provides two model specification tests. Section 3 presents an extensive Monte Carlo simulation study and Sect. 4 covers our empirical application. Finally, Sect. 5 concludes. The Supplementary Material contains all proofs of our theoretical results, as well as the results of the extensive MC study.

2 A cross sectional correlation based specification test for a system of SAR(m) models

In this section, we initially introduce a system of SAR(m) models of order m with $m \in \mathbb{N}$ and discuss its assumptions and efficient parameter estimation procedures. We then present two model specification tests, with the latter showing better power properties as demonstrated in Sect. 3.

2.1 SAR(m): assumptions and parameter estimations

We consider observations y_{it} for individuals $i = 1, \dots, n$ and time periods $t = 1, \dots, T$. For each t , the y_{it} can be modeled by a spatially autoregressive model of order $m \in \mathbb{N}$, SAR(m), i.e.

$$\mathbf{y}_t = \sum_{i=1}^m \rho_i W_i \mathbf{y}_t + \boldsymbol{\varepsilon}_t \quad (1)$$

where $\mathbf{y}'_t = (y_{t1}, \dots, y_{tn})$ and $\boldsymbol{\varepsilon}'_t = (\varepsilon_{t1}, \dots, \varepsilon_{tn})$. The parameters ρ_i with $i = 1, \dots, m$ are the scalar spatial autoregressive coefficients and W_i for $i = 1, \dots, m$ the pre-specified $n \times n$ spatial weighting matrices that do not vary over time. An overview of commonly used spatial matrices is given in Piras et al. (2012). By rearranging, we obtain a simplified version of model (1), i.e.

$$\mathbf{y}_t = (I_n - \sum_{i=1}^m \rho_i W_i)^{-1} \boldsymbol{\varepsilon}_t. \quad (2)$$

The observations in the cross-sectional dimension i are assumed to be fixed. We impose the following model assumptions:

- Assumption 1**
1. The sequence $\{\boldsymbol{\varepsilon}_t\}_{t \in \mathbb{N}}$ has zero mean, is stationary and ergodic.
 2. For $i \in \{1, \dots, m\}$, $r = 1, \dots, n$, $s = 1, \dots, n$, $W_{i,rs} \geq 0$, $W_{i,rr} = 0$.
 3. For $i \in \{1, \dots, m\}$ and $r = 1, \dots, n$, $\sum_{s=1}^n W_{i,rs} = 1$.
 4. The parameter space \mathcal{S}_ρ is defined as $\mathcal{S}_\rho = \{\boldsymbol{\rho} \in \mathbb{R}^m : \|\boldsymbol{\rho}\|_1 < 1\}$ where $\|\cdot\|_1$ defines the L^1 -norm, i.e. $\|\boldsymbol{\rho}\|_1 = \sum_{i=1}^m |\rho_i|$.
 5. The spatial parameter vector $\boldsymbol{\rho}$ is uniquely identified.
 6. Define $\Sigma \stackrel{d}{=} \text{Cov}(\boldsymbol{\varepsilon}_t)$. For $t \in \mathbb{Z}$, $\text{diag}(\Sigma) = (\sigma_1^2, \dots, \sigma_n^2) \in \mathbb{R}^n$.
 7. Each element of the vector $\left(\frac{1}{\sqrt{T}} \sum_{t=1}^T \boldsymbol{\varepsilon}_t \boldsymbol{\varepsilon}'_t\right)_{i < j}$ meets the assumption of a central limit theorem and the corresponding long-term covariances

$$\sum_{s,t \in \mathbb{N}} \text{Cov}[\boldsymbol{\varepsilon}_{it} \boldsymbol{\varepsilon}_{jt}, \boldsymbol{\varepsilon}_{ks} \boldsymbol{\varepsilon}_{ls}]$$

are finite for every $i < j$ and $k < l$, where we interpret $(\cdot)_{i < j}$ as the stacked vector of the upper triangular matrix.

The zero mean and stationarity condition in Assumption 1.1. are plausible especially in the context of daily stock returns (see Aue et al. 2009). If this assumption is violated, trend adjustment, centering or, to be more general, considering appropriate residuals, could ensure that it is (asymptotically) met. To exclude self-neighbors, the diagonal elements of W_i with $i = 1, \dots, m$ are conventionally set equal to zero (Assumption 1.2.). Additionally, Assumption 1.2. claims that all elements are non-negative (which is natural), as distances are measured. Assumption 1.3. ensures that the matrices are bounded and standardized. Assumption 1.4. restricts the parameter space to the sum of the absolute values of the elements of $\boldsymbol{\rho} \in \mathbb{R}^m$ smaller than 1. While the assumption could slightly be generalized (cf. Piras et al. 2012), we follow the notation of Arnold et al. (2013) as it guarantees that the matrix $(I_n - \sum_{i=1}^m \rho_i W_i)$ is

non-singular.¹ Assumption 1.5. yields a high-level identification assumption which is specified in the Supplementary Material in Assumption 3.1. It rules out certain combinations of spatial weighting matrices, e.g., these matrices must be pairwise distinct. Assumption 1.6. allows for heteroscedastic error variances and Assumption 1.7. is a standard asymptotic condition that allows for deriving non-degenerate asymptotic distributions of the estimators and test statistics.

We consider the two step GMM procedure of Arnold et al. (2013) in order to obtain efficient estimates for the parameters ρ_i with $i = 1, \dots, m$ and σ_l^2 with $l = 1, \dots, n$.

This consists of the following two steps:

First, we estimate the spatial parameters $\rho_i, i = 1, \dots, m$ using the method of moments. Due to its construction, this step does not depend on the parameters of variance σ_l^2 with $l = 1, \dots, n$. Second, for given spatial estimates, the estimation of variance parameters is obtained by computing the mean of the estimated $\hat{\varepsilon}_{it}^2$ with $i = 1, \dots, n$. Under some regularity assumptions, the GMM estimator $\hat{\rho}$ is consistent and asymptotically normal (cf. Theorem 2.2). While this is worked out in Arnold et al. (2013) for the special case of $m = 3$, a detailed derivation for the GMM estimator in the general case is presented in the Supplementary Material 2.2, i.e. $m \in \mathbb{N}$ is finite and fixed.

2.2 A two step GMM estimation procedure for a system of SAR(m) models

In the following, we assume that Assumption 1 is fulfilled. The covariance matrix of $\mathbf{u}_t = (I_n - \sum_{i=1}^m \rho_i W_i)^{-1} \boldsymbol{\varepsilon}_t$ is given by

$$\text{Cov}[\mathbf{u}_t] = \left(I_n - \sum_{i=1}^m \rho_i W_i \right)^{-1} \Sigma \left(I_n - \sum_{i=1}^m \rho_i W_i' \right)^{-1} =: V.$$

For the estimation, a two step procedure is considered: First, we estimate the correlation parameters by the method of moments which does not depend on the parameters of variance. Secondly, we estimate the variance parameters.

The moment estimator for the correlation parameters uses the following m moment conditions:

$$E \left[\boldsymbol{\varepsilon}_t' W_i \boldsymbol{\varepsilon}_t \right] = \text{tr}(W_i \Sigma) = 0 \quad \text{for} \quad i = 1, \dots, m. \tag{3}$$

Clearly, the variance parameters σ_i^2 for $i = 1, \dots, m$ do not enter the moment conditions. Replacing $\boldsymbol{\varepsilon}_t$ by

$$\boldsymbol{\varepsilon}_t = \left(I_n - \sum_{i=1}^m \rho_i W_i \right) \mathbf{u}_t$$

¹ The matrix $(I_n - \sum_{i=1}^m \rho_i W_i)$ is strictly diagonally dominant.

and averaging over t gives the theoretical system of equations

$$\Gamma \boldsymbol{\lambda} + \boldsymbol{\gamma} = 0,$$

where $\boldsymbol{\lambda} = \boldsymbol{\lambda}(\boldsymbol{\rho})$ is a functional vector of $\boldsymbol{\rho} = (\rho_1, \dots, \rho_m)$ of dimension $M = \binom{m}{1} + \binom{m+2-1}{2}$, $\binom{\cdot}{\cdot}$ denoting the binomial coefficient, such that

$$\lambda_i = \rho_i \quad \text{for } i = 1, \dots, m \quad (4)$$

$$\lambda_{m+i} = \rho_i^2 \quad \text{for } i = 1, \dots, m \quad (5)$$

$$\lambda_{2m+\#\{ij \mid i < j, i < l, j \leq k\}} = \rho_l \rho_k \quad \text{for } l, k = 1, \dots, m, \quad (6)$$

where $\#\{ij \mid i < j, i < l, j \leq k\}$ represents the number of integer pairs ij such that the conditions $i < j, i < l$ and $j \leq k$ are fulfilled for $l, k = 1, \dots, m$. The elements of $\Gamma \in \mathbb{R}^{m \times M}$ and $\boldsymbol{\gamma} \in \mathbb{R}^m$ are defined by for $i, j = 1, \dots, m$,

$$\Gamma_{i,j} = \mathbb{E} \left[-\frac{1}{T} \sum_{t=1}^T \mathbf{u}'_t (W_i + W'_i) W_j \mathbf{u}_t \right], \quad (7)$$

$$\Gamma_{i,m+j} = \mathbb{E} \left[\frac{1}{T} \sum_{t=1}^T \mathbf{u}'_t W'_j W_i W_j \mathbf{u}_t \right], \quad (8)$$

$$\Gamma_{i,2m+\#\{ij \mid i < j, i < l, j \leq k\}} = \mathbb{E} \left[\frac{1}{T} \sum_{t=1}^T \mathbf{u}'_t W'_l (W_i + W'_i) W_k \mathbf{u}_t \right],$$

$$\gamma_i = \mathbb{E} \left[\frac{1}{T} \sum_{t=1}^T \mathbf{u}'_t W_i \mathbf{u}_t \right]. \quad (9)$$

Let G and \mathbf{g} be the empirical counterparts of Γ and $\boldsymbol{\gamma}$, i.e. the expectation operator is left out. The moment estimator for $\boldsymbol{\rho} = (\rho_1, \dots, \rho_m)'$ is defined as

$$\hat{\boldsymbol{\rho}} = (\hat{\rho}_1, \dots, \hat{\rho}_m)' = \arg \min_{\boldsymbol{\rho} \in S} \|G\boldsymbol{\lambda} + \mathbf{g}\|$$

where $\|\cdot\|$ represents the euclidean norm.

Remark 2.1 For $k, l \in \{1, \dots, m\}$, the entries of $\mathbb{E}[G] = \Gamma$ given in (7)–(9) can be calculated as

$$\Gamma_{k,l} = \text{tr}((W_k + W'_k) W_l V),$$

$$\Gamma_{k,m+l} = \text{tr}(W'_l W_k W_l V),$$

$$\Gamma_{i,2m+\#\{ij \mid i < j, i < l, j \leq k\}} = \text{tr}(W'_l (W_i + W'_i) W_k V).$$

Since the theoretical term $\Gamma \boldsymbol{\lambda} + \boldsymbol{\gamma}$ is equal to zero for the true parameter values, the moment estimator for $\hat{\boldsymbol{\rho}}$ minimizes the corresponding empirical system $G\boldsymbol{\lambda} + \mathbf{g}$. Arnold et al. (2013) prove consistency and asymptotic normality of the moment estimator (cf. Theorem 2.2) for $T \rightarrow \infty$, for which an additional assumption is needed.

Assumption 2 1. The true parameter $\rho \in S$ is the unique solution of the theoretical system of equations, i.e.

$$\Gamma\lambda + \gamma = 0 \Leftrightarrow \hat{\rho} = \rho.$$

2. The matrix $E\left(\frac{\partial(G\lambda+g)}{\partial\hat{\rho}}(\mathbf{y}_t, \rho)\right) =: \mathbf{d} = \Gamma\lambda^{(1)}$ exists, is finite and has full rank with $\lambda^{(1)}$ a $(M \times m)$ dimensional matrix defined as

$$\begin{aligned} \lambda^{(1)}(l, l) &= 1, & \lambda^{(1)}(2m + \#\{ij \mid i < j, i < l, j \leq k\}, l) &= \rho_k \\ \lambda^{(1)}(m + l, l) &= 2\rho_l, & \lambda^{(1)}(2m + \#\{ij \mid i < j, i < l, j \leq k\}, k) &= \rho_l \end{aligned}$$

for all $l, k = 1, \dots, m$.

3. For

$$f(\mathbf{u}_t, \rho) = \begin{pmatrix} \boldsymbol{\varepsilon}'_t W_1 \boldsymbol{\varepsilon}_t \\ \vdots \\ \boldsymbol{\varepsilon}'_t W_m \boldsymbol{\varepsilon}_t \end{pmatrix},$$

it holds that, for $j \rightarrow \infty$, $E[f(\mathbf{u}_t, \rho) \mid f(\mathbf{u}_{t-j}, \rho), f(\mathbf{u}_{t-j-1}, \rho), \dots]$ converges in mean square to zero and that, for

$$\begin{aligned} \mathbf{v}_j &= E[f(\mathbf{u}_t, \rho) \mid f(\mathbf{u}_{t-j}, \rho), f(\mathbf{u}_{t-j-1}, \rho), \dots] \\ &\quad - E[f(\mathbf{u}_t, \rho) \mid f(\mathbf{u}_{t-j-1}, \rho), f(\mathbf{u}_{t-j-2}, \rho), \dots] \end{aligned}$$

the infinite sum $\sum_{t=-\infty}^{\infty} E[(\mathbf{v}_j \mathbf{v}_j)^\frac{1}{2}]$ is finite.

Under the Assumptions 1 and 2 the GMM estimator $\hat{\rho}$ is consistent and asymptotic normal as the following theorem shows:

Theorem 2.2 *Let Assumptions 1 and 2 hold. Then, for $S_W = \sum_{t=-\infty}^{\infty} E[f(\mathbf{u}_t, \rho) f(\mathbf{u}_t, \rho)']$ and $T \rightarrow \infty$ it holds:*

1. $\hat{\rho} \xrightarrow{p} \rho$
2. $\sqrt{T}(\hat{\rho} - \rho) \xrightarrow{d} N(0, \mathbf{d}^{-1} S_W (\mathbf{d}^{-1})')$.

The proof of this theorem is similar to the proof of Theorem 2.1 in Arnold et al. (2013).

2.3 The specification test

Our proposed test is a variation of the classical Portmaneau test, i.e. we check if the covariance matrix of the idiosyncratic error resembles a diagonal matrix. Under the null hypothesis, the off-diagonal elements do not deviate too far from zero. Let $\hat{H} \in \mathbb{R}^{n \times n}$ denote the empirical covariance matrix of the residuals times the square root of the

time horizon T , i.e. $\hat{H} = \sqrt{T} \widehat{\text{Cov}} [\hat{\boldsymbol{\varepsilon}}_t]$ and \hat{H}_{ij} its elements with $i, j \in \{1, 2, \dots, n\}$. Let σ_{ij}^2 denote the (i, j) -th element of the theoretical counterpart Σ , i.e. the error covariance matrix. Since \hat{H} and Σ are symmetric, it is sufficient to consider only the elements of the upper triangle of the matrix Σ . Hence, the null hypothesis is given by

$$H_0 : \sigma_{ij}^2 = 0 \text{ for all } i < j \quad \text{vs.} \quad H_1 : \exists i, j \text{ with } i < j : \sigma_{ij}^2 \neq 0. \quad (10)$$

We opt to use χ^2 -type tests for this testing problem. Instead of considering each element or the maximum of the absolute value of all off-diagonals, we take the sum of each element squared into account. Thus, the test statistic is given by

$$S = \sum_{i < j, i, j = 1, \dots, n} (\hat{H}_{ij})^2. \quad (11)$$

The aim of the following theorem is to decompose the limit in distribution of the empirical covariance matrix times \sqrt{T} into three matrices, which allow to determine the limit in distribution of the sum of the elements of the upper triangular matrix. Here and in the following $\underset{T \rightarrow \infty}{\text{dlim}}$ denotes limit in distribution and $\stackrel{d}{=}$ equality in distribution.

Theorem 2.3 *Under the null hypothesis $H_0 : \sigma_{ij}^2 = 0$ for all $i < j$ and the assumptions of Theorem 2.2, the following holds for $1 \leq i, j \leq n$*

$$\underset{T \rightarrow \infty}{\text{dlim}} \sqrt{T} \widehat{\text{Cov}} [\hat{\boldsymbol{\varepsilon}}_t] = A + B + B' \in \mathbb{R}^{n \times n} \quad (12)$$

with $A_{ii} = \lim_{T \rightarrow \infty} \sqrt{T} \sum_{t=1}^T \sigma_{it}^2 = \infty$ and $A_{ij} \sim N \left(0, \lim_{T \rightarrow \infty} \text{Var} \left[\frac{1}{\sqrt{T}} \sum_{t=1}^T \boldsymbol{\varepsilon}_{it} \boldsymbol{\varepsilon}_{jt}' \right] \right)$,

where

$\text{Cov}[A_{ij}, A_{kl}] = 0$ for $i \neq j$ and $k \neq l$ with $(i, j) \neq (k, l)$. We also define

$$B \stackrel{d}{=} \left(\sum_{i=1}^m X_i W_i \right) \left(I_n - \sum_{i=1}^m \rho_i W_i \right)^{-1} \Sigma, \text{ where}$$

$$\mathbf{X} \stackrel{d}{=} (X_1, \dots, X_m) \sim N(0, \mathbf{d}^{-1} S_W (\mathbf{d}^{-1})') \in \mathbb{R}^{1 \times m}, \quad S_W \stackrel{d}{=} \sum_{t=-\infty}^{\infty} \text{E}[f(\mathbf{u}_t, \boldsymbol{\rho}) f(\mathbf{u}_t, \boldsymbol{\rho})']$$

with $f(\mathbf{u}_t, \boldsymbol{\rho}) \stackrel{d}{=} (\boldsymbol{\varepsilon}'_t W_1 \boldsymbol{\varepsilon}_t, \dots, \boldsymbol{\varepsilon}'_t W_m \boldsymbol{\varepsilon}_t)'$ and \mathbf{d} defined in Assumption 2.

Three remarks about Theorem 2.3 are in order. First, the leading elements of matrix A diverge to infinity. However, the tests consider only the off-diagonal elements ($i \neq j, i, j = 1, \dots, n$) which are finite by Assumption 1.7. This in turn ensures that the test is well defined. Secondly, since $(I_n - \sum_{i=1}^m \rho_i W_i)$ is strictly diagonally dominant, the inverse exists. Thirdly, we note that the matrices B and its transposed appear in the limit. This is due to the fact of estimating $\boldsymbol{\rho}$ instead of using the unknown population quantity. The analysis of such a residual effect (see Demetrescu and Wied 2019) is

somewhat complicated, since the additional terms need different standardizing factors in the proof.² However, all terms in the limiting distribution are based on the same error terms. Thus, the convergence is jointly and the limiting distribution in (12) is multivariate normal. If we additionally assume serial independence in the error vector, the variance of the elements in the limiting matrix A simplifies to a product shown in the following remark.

Remark 2.4 Suppose the assumptions of Theorem 2.3 hold. If $\{\boldsymbol{\varepsilon}_t\}_{t \in \{1, \dots, T\}}$ is serially independent, then

$$A_{ij} \sim N(0, \sigma_i^2 \sigma_j^2) \text{ for } i \neq j. \tag{13}$$

In accordance with our test statistic S (11), we can reformulate the test in vectorial notation, i.e.

$$S = \hat{\boldsymbol{\alpha}}' \hat{\boldsymbol{\alpha}}, \tag{14}$$

where $\hat{\boldsymbol{\alpha}}$ represents the vector of the upper triangle of the empirical covariance matrix of the residuals times \sqrt{T} , i.e. \hat{H} . Since the empirical covariance matrix consists of n^2 elements, the upper triangular matrix vector consists of $n(n - 1)/2$ elements and has the following form:

$$\begin{aligned} \hat{\boldsymbol{\alpha}} &\stackrel{d}{=} \text{dlim}_{T \rightarrow \infty} \left(\sqrt{T} \widehat{\text{Cov}} [\hat{\boldsymbol{\varepsilon}}_t] \right)_{i < j, i, j = 1, \dots, n} \\ &= \text{dlim}_{T \rightarrow \infty} \left(\frac{1}{\sqrt{T}} \sum_{i < j, i, j = 1, \dots, n} \hat{\boldsymbol{\varepsilon}}_t \hat{\boldsymbol{\varepsilon}}_t' \right) = \text{dlim}_{T \rightarrow \infty} \frac{1}{\sqrt{T}} \sum_{t=1}^T \hat{\boldsymbol{\varepsilon}}_t^* \in \mathbb{R}^{\frac{n(n-1)}{2}} \\ &\text{with } \hat{\boldsymbol{\varepsilon}}_t^* \stackrel{d}{=} (\hat{\varepsilon}_{1t} \hat{\varepsilon}_{2t}, \dots, \hat{\varepsilon}_{1t} \hat{\varepsilon}_{nt}, \hat{\varepsilon}_{2t} \hat{\varepsilon}_{3t}, \dots, \hat{\varepsilon}_{2t} \hat{\varepsilon}_{nt}, \dots, \hat{\varepsilon}_{(n-1)t} \hat{\varepsilon}_{nt})'. \end{aligned}$$

By means of Slutsky’s theorem we define the theoretical counterpart

$$\begin{aligned} \boldsymbol{\alpha} &\stackrel{d}{=} (A)_{i < j, i, j = 1, \dots, n} \\ &= \text{dlim}_{T \rightarrow \infty} \left(\frac{1}{\sqrt{T}} \sum_{i < j, i, j = 1, \dots, n} \boldsymbol{\varepsilon}_t \boldsymbol{\varepsilon}_t' \right) = \text{dlim}_{T \rightarrow \infty} \frac{1}{\sqrt{T}} \sum_{t=1}^T \boldsymbol{\varepsilon}_t^* \in \mathbb{R}^{\frac{n(n-1)}{2}} \\ &\text{with } \boldsymbol{\varepsilon}_t^* \stackrel{d}{=} (\varepsilon_{1t} \varepsilon_{2t}, \dots, \varepsilon_{1t} \varepsilon_{nt}, \varepsilon_{2t} \varepsilon_{3t}, \dots, \varepsilon_{2t} \varepsilon_{nt}, \dots, \varepsilon_{(n-1)t} \varepsilon_{nt})' \end{aligned}$$

which stacks the upper triangular matrix of the covariance matrix of the errors times \sqrt{T} in a vector. Analogously, $\boldsymbol{\delta}$ defines the vector of the stacked upper triangular matrix of B and $\boldsymbol{\delta}^*$ of B' , respectively, i.e. for $Z_W = \text{dlim}_{T \rightarrow \infty} \sum_{g=1}^m \sqrt{T} (\rho_g - \hat{\rho}_g) W_G$ we define

² For a detailed analysis of the convergence rate we refer to Lemma A.1 in the Supplementary Material.

$$\boldsymbol{\delta} \stackrel{d}{=} (B)_{i < j, i, j=1, \dots, n} = \left(Z_W (I_n - \sum_{g=1}^m \rho_g W_G)^{-1} \Sigma \right)_{i < j, i, j=1, \dots, n} \in \mathbb{R}^{\frac{n(n-1)}{2}},$$

$$\boldsymbol{\delta}^* \stackrel{d}{=} (B')_{i < j, i, j=1, \dots, n} = \left(\Sigma' (I_n - \sum_{g=1}^m \rho_g W_G')^{-1} Z_W' \right)_{i < j, i, j=1, \dots, n} \in \mathbb{R}^{\frac{n(n-1)}{2}}.$$

The vectors $\boldsymbol{\delta}$ and $\boldsymbol{\delta}^*$ are well defined, since B is not necessarily symmetric.

Lemma 2.5 $\boldsymbol{\delta}$ represents the stacked vector of the upper triangle and $\boldsymbol{\delta}^*$ of the lower triangle of the matrix $Z_W (I_n - \sum_{g=1}^m \rho_g W_G)^{-1} \Sigma$, i.e. for $i, j \in \{1, \dots, n\}$

$$\boldsymbol{\delta}^* = \left(Z_W (I_n - \sum_{g=1}^m \rho_g W_G)^{-1} \Sigma \right)_{i > j, i, j=1, \dots, n} \in \mathbb{R}^{\frac{n(n-1)}{2}}. \quad (15)$$

The next lemma provides the limit distribution of our test statistic S (14).

Lemma 2.6 Suppose the assumptions of Theorem 2.3 hold. Then the test statistic S (11) is asymptotically distributed as

$$S = \hat{\boldsymbol{\alpha}}' \hat{\boldsymbol{\alpha}} \xrightarrow[T \rightarrow \infty]{d} (\boldsymbol{\alpha} + \boldsymbol{\delta} + \boldsymbol{\delta}^*)' (\boldsymbol{\alpha} + \boldsymbol{\delta} + \boldsymbol{\delta}^*),$$

where the covariance matrix for $\boldsymbol{\alpha}$ is given by

$$\text{Cov} [\boldsymbol{\alpha}] = \begin{pmatrix} \lim_{T \rightarrow \infty} \text{Var} \left[\frac{1}{\sqrt{T}} \sum_{t=1}^T \boldsymbol{\varepsilon}_{1t} \boldsymbol{\varepsilon}_{2t}' \right] \cdots & & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lim_{T \rightarrow \infty} \text{Var} \left[\frac{1}{\sqrt{T}} \sum_{t=1}^T \boldsymbol{\varepsilon}_{(n-1)t} \boldsymbol{\varepsilon}_{nt}' \right] \end{pmatrix}.$$

To obtain the corresponding critical values, first, the model parameters are estimated for all models considered in the empirical application below. Second, a bootstrap procedure is employed to generate the critical values by drawing from the estimated model. Tables 6 and 7 show the corresponding critical values for all models considered in the empirical application below. As shown in Sect. 3, our proposed test takes care of size demands and has good power properties. The next subsection presents a modification of the proposed specification test with a simpler limit distribution.

2.4 Simplified tests

In Theorem 2.3 we have shown that the elements of the limiting distribution follow a multivariate normal distribution. Thus, if we standardize the test statistic S (14) by its covariance matrix, we get a new test statistic S_χ^* which is χ^2 -distributed, i.e.

$$S_{\chi}^* = \hat{\alpha}' (\text{Cov} [\alpha + \delta + \delta^*])^{-1} \hat{\alpha} \sim \chi_{\frac{n(n-1)}{2}}^2. \tag{16}$$

The quantiles of this limit distributions can be easily obtained, but the implementation of test statistic is more complicated than that of (11). The following discussion shows how the implementation can be simplified.

The terms δ and δ^* can be regarded as additional noise which comes from the estimation procedure. This additional noise can be extracted by decomposing the covariance matrix given in (16) into two parts. Thus, we have

$$\text{Cov} [\alpha + \delta + \delta^*] = \text{Cov} [\alpha] + \Psi \tag{17}$$

with $\Psi = \text{Cov} [\delta + \delta^*] + \text{Cov} [\alpha, \delta + \delta^*] + \text{Cov} [\alpha, \delta + \delta^*]'$. The first part $\text{Cov} [\alpha]$ covers the underlying variance structure while the second part Ψ can be considered as a noise term.³

If either $\|(\text{Cov} [\alpha])^{-1} \Psi\| < 1$ or $\|\Psi(\text{Cov} [\alpha])^{-1}\| < 1$ holds,⁴ the inverse of the covariance matrix (17) can be estimated by means of a Taylor series approximation and a telescoping sum.⁵ This yields

$$\begin{aligned} (\text{Cov} [\alpha + \delta + \delta^*])^{-1} &= (\text{Cov} [\alpha])^{-1} - (\text{Cov} [\alpha])^{-1} \Psi (\text{Cov} [\alpha])^{-1} \\ &\quad + (\text{Cov} [\alpha])^{-1} \Psi (\text{Cov} [\alpha])^{-1} \Psi (\text{Cov} [\alpha])^{-1} - \dots \\ &\leq (\text{Cov} [\alpha])^{-1}. \end{aligned}$$

Thus, $(\text{Cov} [\alpha])^{-1}$ is an upper bound for the inverse of the covariance matrix (17). Hence, for

$$S_{\chi} = \lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=1}^T \hat{\epsilon}_t^{*'} (\text{Cov} [\alpha])^{-1} \sum_{t=1}^T \hat{\epsilon}_t^*, \tag{18}$$

we have the algebraic relation $S_{\chi} \geq S_{\chi}^* \sim \chi_{\frac{n(n-1)}{2}}^2$. Therefore, using the test statistic S_{χ} with the quantiles of the $\chi_{\frac{n(n-1)}{2}}^2$ leads to a *liberal* test. This test might be preferred in applications if it appears to be too time-consuming to generate critical values by simulations. In order to study the behavior of S and S_{χ} in finite samples, we perform an extensive Monte Carlo simulation study which can be found in the next section.

³ If we additionally assume serial independence, the covariance matrix of α can easily be implemented, since only the variances need to be estimated, cf. Lemma A.3. Otherwise, the covariance matrix of α is given in Lemma 2.6.

⁴ In our Monte Carlo simulation we observed that this is usually the case whenever the variance of ϵ_{it} is greater than 1 for all $i = 1, \dots, n$.

⁵ The sum and product of two symmetric positive semidefinite (psd) matrices is still psd.

3 Monte Carlo simulation

3.1 Serial independence

The Monte Carlo (MC) simulation study consists of three main simulations, and in all simulations we assume a SAR(m) for $m = 3, 4$. The results of our extensive MC simulation study are provided in the Supplementary Material. While the first two simulations assume serial independence, the third simulation examines the behavior of the test in the case of GARCH(1,1) driven errors. To examine size and power properties of the test S (2.3), we draw $B = 300$ times from the asymptotic limit distribution given in Lemma (2.6). The overall number of MC repetitions is equal to 701.

FIRST SETTING: The first simulation depicts a SAR(3) model, i.e.

$$\mathbf{y}_t = \rho_1 W_1 \mathbf{y}_t + \rho_2 W_2 \mathbf{y}_t + \rho_3 W_3 \mathbf{y}_t + \boldsymbol{\varepsilon}_t, \quad (19)$$

where $\mathbf{y}_t, \boldsymbol{\varepsilon}_t \in \mathbb{R}^n$ for $t = 1, \dots, T$ and $(W_1)_{ij} = \frac{1}{n-1}$ for all $i \neq j$ and $(W_1)_{ii} = 0$. The spatial matrices W_2 and W_3 are defined as

$$(W_2)_{ij} \stackrel{d}{=} \begin{cases} 1, & \text{if } j \text{ even and } i \neq j \\ 1, & \text{if } j - 1 = i \\ 0, & \text{otherwise} \end{cases}$$

$$(W_3)_{ij} \stackrel{d}{=} \begin{cases} 1/(n/2 - 1), & \text{if } i, j \leq \frac{n}{2} \text{ or } i, j > \frac{n}{2} \\ 0, & \text{otherwise,} \end{cases}$$

where additionally the matrix W_2 is row standardized by its row sum $\sum_j (W_2)_{ij}$ for $i = 1, \dots, n$. In terms of interpretation, the matrix W_1 can be regarded as a weighting matrix, where each firm has the same weight. Thus, the matrix W_1 captures a general effect, e.g. global crisis, market performance in the past etc.⁶ Another interpretation in terms of networks might be that there is a fully connected network, i.e. that all stocks are related to each other in the same manner. The spatial matrix W_2 is an example of an asymmetric weighting matrix. This might be interesting from an economic point of view, if there is evidence that a particular (large) country has much influence on another (smaller) country, but that there is less dependence in the other direction.

In economic analyses, W_3 functions as a pivotal dichotomous matrix, segmenting the market into distinct sectors. This dichotomy underscores the contrast between entities who benefit from specific interventions, e.g. tax reforms, aid disbursements, or market shifts, and those who remain unaffected or neutral to these changes. This dichotomous framework is particularly apt for policy evaluation, as it offers a nuanced insight into the differential impacts and subsequent redistributive consequences of policies or market dynamics.

⁶ Even if W_1 is equally weighted, ρ_1 cannot be considered as a fixed affect which affects market participant equally, since fixed effects are time independent. SAR models try to capture this time dependence structure with fixed weighting matrices.

The vector of observation \mathbf{y}_t is generated by a multivariate normal error vector $\boldsymbol{\varepsilon}_t$ with zero mean and covariance matrix $\Sigma = \sigma^2 I_n$, where I_n represents the n -dimensional identity matrix and the model representation in (19). The parameter of spatial dependence is given by $\boldsymbol{\rho} = (0.45, 0.3, 0.15)$ and the homoscedastic variance equals $\sigma^2 = 2$.

To calculate the power of our tests, we do not simulate the errors from a multivariate normal with a diagonal covariance matrix. Instead, we use the following misspecification: If we consider a market with n participants, then there are $n(n-1)/2$ possible pairs that represent the off-diagonal elements of the covariance matrix. After a simple transformation these off-diagonal elements can be considered as participants that are correlated with each other. The parameter ζ describes the portion of pairs,⁷ the parameter κ^2 describes their correlation. E.g. if we consider a market that consists of $n = 20$ actors, then there are $n(n-1)/2 = 190$ different pairs. If $\zeta = 0.1$ and $\kappa = 0.2$, we assume that there are 19 pairs that have a correlation coefficient that is equal to 0.04. No further assumptions are made about the structure of correlation. However, the correlation structure is completely random,⁸ i.e. we predetermine only generally the proportion of correlated pairs and their correlation and not the correlation of specific pairs. Results are presented in Table 1 in the Supplementary Material.

RESULTS FOR S : Collectively, the test holds the size level. In small samples, i.e. whenever the ratio of T over n is small ($T \leq 100$, $n \leq 50$) and the dependence structure in the error term is more or less negligible (cf. $\kappa = \zeta = 0.05$) the power of the test is low. However, if there are sufficient observations (i.e. $\frac{T}{n} > 10$) and if the dependence structure in the data set is not negligible ($\kappa, \zeta \geq 0.1$), the test provides good power properties even in small samples. All in all we observe an increasing power whenever the dependence structure (κ or ζ) or the number of observations (n or T) increase.

RESULTS FOR S_χ : Similar results are obtained for the second test S_χ (11) which can be found in Table 2.⁹ In small samples, S_χ performs worse than S in terms of size and power. This is due to the fact that we are using the empirical approximation for the inverse covariance matrix that is employed in S_χ , which is biased in small samples. Consequently, as T tends to infinity the size of the test S_χ converges to the desired nominal level of 5% and the power increases as the level of misspecification rises. However, additional simulations show that the tests' power decreases in the case of too large ζ , i.e. a highly non-sparse covariance matrix. Here, the population moment conditions of the GMM estimate (cf. equation (3) given in the Supplementary Material) are severely violated so that the model is misspecified and the behavior of the model estimators $\hat{\rho}$ is unclear (Fleming 2004).

To summarize, both tests show good size and power properties whenever the ratio T over n is greater than 10. Based on the simple limiting distribution of S_χ^* , the test S_χ is also very easy to implement since the test statistic S_χ requires only the empirical covariance matrix of the residuals.

⁷ In case that $\zeta \cdot n(n-1)/2$ is odd we round down.

⁸ This procedure of misspecification ensures that the moment conditions (3) are violated, thus, the GMM estimator is biased (Hansen 1982).

⁹ The second test is applicable since for every simulation it holds true that either $\|(\text{Cov}[\boldsymbol{\alpha}])^{-1}\Psi\| < 1$ or $\|\Psi(\text{Cov}[\boldsymbol{\alpha}])^{-1}\| < 1$.

SECOND SETTING: For the second MC simulation we consider a SAR(4) model

$$\mathbf{y}_t = \rho_1 W_1 \mathbf{y}_t + \rho_2 W_2 \mathbf{y}_t + \rho_3 W_3 \mathbf{y}_t + \rho_4 W_4 \mathbf{y}_t + \boldsymbol{\varepsilon}_t, \quad t = 1, \dots, T,$$

where W_1 is a group interaction matrix of the first two-thirds (the off-diagonal elements of the first two-third upper sub-matrix are set to $1/(\frac{2}{3}n - 1)$ and all other elements to zero), W_2 is a group interaction matrix of the last third (cf. W_1), W_3 a binary contiguity matrix of the third-order neighbors assuming the observations $1, \dots, n$ are arranged in a circular pattern, e.g., 2 is a third-order neighbor of $n - 1, n, 1$ and $3, 4, 5$, and¹⁰

$$(W_4)_{ij} \stackrel{d}{=} \begin{cases} \frac{1}{2 \cdot \lfloor n-1 \rfloor}, & \text{if } i \text{ is even and } j \text{ odd or vice versa} \\ 0, & \text{otherwise.} \end{cases}$$

The vector of autoregressive parameters ρ is given by $\boldsymbol{\rho} = (-0.2 \ 0.05 \ 0.1 \ 0.5)$. Moreover, we presuppose heteroscedastic normal error terms, i.e. $\sigma_i \sim N(0, 1)$ for $i = 1, \dots, n$. In order to analyze the power in case of misspecification, we choose ζ and κ likewise to the first MC simulation. To determine the size and power we draw $B = 300$ times from the asymptotic limit distribution given in Lemma (2.6). The overall number of MC repetitions is equal to 701. The results of the tests can be found in Table 3.

RESULTS FOR S AND S_χ : Even if the results of the second analysis are not entirely comparable with those from the first simulation,¹¹ it is clearly observable that the tests S and S_χ hold the size level. The power increases if either the correlation structure (κ or ζ) or the number of observation increases (n or T). Thus, the results presented in the second, more complex study are in line with those given in the first simulation.

The next section shows that the test holds size and power demands even if the error terms follow a GARCH process.

3.2 GARCH(1,1)

One of many problems researchers and practitioners face when analyzing financial data is its volatile structure. Volatility of financial data has been extensively studied in the last twenty years. An important aspect of the analysis is volatility clustering, where conditional heteroskedasticity, which leads to an increase in the probability of rare events, can be modeled with GARCH errors. Since the SAR(m) model is a powerful instrument in modeling financial data¹², the third Monte Carlo simulation for our proposed test statistic (14) assumes that the errors of the data generating process (DGP) are driven by a GARCH(1,1) model, i.e. for $t = 1, \dots, T$ and $i = 1, \dots, n$

$$y_{it} = \sigma_{it}(I_n - \rho_1 W_1 - \rho_2 W_2 - \rho_3 W_3)^{-1} \epsilon_{it},$$

¹⁰ Matrices W_1, W_2, W_3 are the counterparts to the matrices $G1, G2, BC3$ given in Piras et al. (2012).

¹¹ The model presupposes heteroscedasticity and the spatial structure is completely different. From this it follows that the violation of the moment condition (3) is not one-to-one comparable.

¹² The empirical analysis in Sect. 4 shows that a SAR(3) seems reasonable in times of no economic crisis.

$$\sigma_{it}^2 = 0.33 + 0.33\sigma_{i(t-1)}^2 + 0.075y_{i(t-1)}^2,$$

$$\epsilon_{it} \stackrel{i.i.d.}{\sim} N(0, 1).$$

To receive comparable results, the spatial matrices W_1, W_2, W_3 are the same as those of the first MC simulation of Sect. 3. The size and power results are presented in Table 4. At first, it should be noted that the number of observation of a GARCH adjusted data set needs to be significantly higher than a data set with no GARCH adjustment, i.e. $T > 1000$, since for the case of a GARCH adjustment an initial estimate needs to be conducted and a primarily high error of estimation violates the stationarity assumption. However, with a sufficiently large set of observations, the test S (14) also performs well with reference to size and power.

4 Empirical analysis

In the empirical application, we analyze the spatial dependencies of the daily stock returns of the Euro Stoxx 50 members for the period from April 2003 to December 2009 and from January 2018 to December 2020, using the composition of January 2010 and January 2018.¹³ The stock prices are adjusted and transferred to log returns.

To model the spatial relations, we consider four spatial matrices W_G, W_C, W_B and W_I , which are designed to capture the dependencies in different ways. The matrix W_G represents general dependencies¹⁴ while the matrices W_C and W_B depict local and industry-specific dependencies, respectively.¹⁵ Finally, the matrix W_I captures the spatial interaction effect of countries and industries, i.e. we affiliate two stocks that belong to the same country and industry. Note that due to construction, none of the spatial matrices W_G, W_C, W_B and W_I are symmetric.

In our analysis, we examine 15 different SAR models, i.e. we consider SAR models, where the number and composition of spatial matrices varies between the models. In the simplest case, only one spatial matrix constitutes our SAR model, e.g., $\mathbf{y}_t = \rho_G W_G \mathbf{y}_t + \boldsymbol{\epsilon}_t$; in the most complex case, all four spatial matrices are included, i.e., $\mathbf{y}_t = \rho_G W_G \mathbf{y}_t + \rho_C W_C \mathbf{y}_t + \rho_B W_B \mathbf{y}_t + \rho_I W_I \mathbf{y}_t + \boldsymbol{\epsilon}_t$. Thus, for $m \in \{1, 2, 3, 4\}$ the general model for the log stock returns on day $t = 1, \dots, T$ is given by

$$\mathbf{y}_t = \sum_{i=1}^m \rho_i W_i \mathbf{y}_t + \boldsymbol{\epsilon}_t \text{ with } W_I \in \{W_G, W_C, W_B, W_I\} \tag{20}$$

where \mathbf{y}_t is the vector of log stock returns on day t and ρ_i the corresponding spatial parameter. In this model, we can interpret the terms $W_i \mathbf{y}_t$ as specific factors, in which external information (about countries and industries) is included. The term $W_G \mathbf{y}_t$

¹³ For the partitioning of the Euro Stoxx 50 members into branches and countries we refer to Table 5.
¹⁴ The off-diagonal elements of W_G are first set to 1 and then standardized using the market capitalization. Thus, W_G captures impacts that affect all stocks in a similar way, such as past stock market performance.
¹⁵ The construction of the matrices W_C and W_B is similar to the one of the matrix W_G : First, the off-diagonal elements are set to 1 if to stocks belong to the same country (W_C) or industry (W_B), respectively. Finally, the rows are standardized based on the corresponding market capitalization of the associated stocks.

represents the market factor. Our empirical analysis is motivated by two key questions: First, how well can the different SAR models explain the movement of stocks and which spatial matrices do we need for a sufficiently good model? Secondly, can we use the model specification test applied to the 15 different models to ascertain which spatial matrix provides the greatest explanatory power?

To answer these questions, we consider a rolling window of the length of one quarter ($T = 63$), i.e., we check whether our model specification test rejects the null hypothesis that the log returns can be modeled by an SAR(m) model for $m \in \{1, 2, 3, 4\}$. Furthermore, we use the observation period of the rolling window to calculate one-day Value-at-Risk (VaR) forecasts for each model to see if the in-sample performances of the models (as measured by our new test) correspond to the out-of-sample performances. This means that we check if there is a dependence between high proportions of incorrect VaR forecasts and low p-values of the tests. If this dependence exists, our new specification test might be interpreted as a backtest in the spirit of Ziggel et al. (2014) among others. All analyses are performed on a significance level of 5%. The number of draws from the limit distribution is set to $B = 300$. The results for the time periods from 2003 to 2009 and 2018 to 2020 are depicted in Figs. 1 and 2, respectively.

Figure 1 shows the results of our suggested model specification test applied to the 15 models from (20) in a rolling window for the period from April 2003 to December 2009 of the Euro Stoxx 50. In order to apply the model specification test we initially calculate the spatial parameters of the underlying model using a two step GMM estimation procedure (cf. Sect. 2.2) and then apply our model specification test. The blue line in the subplots of Fig. 1 describes the ratio of the test statistic and its 95%-quantile. Thus, if the value is greater than 1, there is no statistical evidence to reject the null hypothesis. A value smaller than 1, however, provides evidence against the underlying model.

In addition, we generate VaR forecasts with standard normally distributed errors for the minimum-variance portfolio based on the underlying model on a rolling window of size $T = 63$. The proportion of incorrect VaR forecasts is represented by the number in each subplot. In Figure 1, a total of four core observations can be made:

First, it seems that more complex models with at least two spatial matrices describe the data better. On the one hand, the period in which the null hypothesis of a correct model assumption cannot be rejected is longer compared to models incorporating only one spatial matrix and on the other hand, false VaR predictions are close to the significance level $\alpha = 5\%$. Secondly, Fig. 1 illustrates that in periods of economic crisis the spatial model (20) is less applicable, since in all models the ratio (blue line) is on a significantly lower level. Particularly, this can be observed in the period of the financial crisis beginning in summer 2007. This is also consistent that in times of bear markets the correlation among market participants increases significantly. Thus, the resulting extensive dependency structure cannot be well captured by a simple SAR(m) model. Accordingly, the results of our test provide evidence that the effects of the dot-com bubble crisis around 2000 last until summer 2004, since the test rejects the application of model (20). In the two following years (2004–2006), however, subplot 6 of Fig. 1 depicts evidence to apply the model, since the blue line is often greater than 1. In the remaining observation period the test indicates that a spatial model is inappropriate,




Fig. 1 Rolling Window for $T = 63$ from 2003 to 2009. Rolling window parameter estimation of size $T = 63$ in a data set of size 1861 and dimension $n = 50$. 15 different models are considered. The first figure describes the SAR(1) model incorporating only a single general dependence structure (W_G). The last figure depicts the results of the SAR(4) model incorporating the spatial matrices W_G, W_C, W_B, W_I . The number of draws from the limit distribution is set to $B = 300$. The blue line depicts the ratio of the 95%-quantile of the limit distribution given in Lemma (2.6) over the test statistic S from (11). A value below 1 indicates that the assumed model is not correct. The orange line is the accumulated spatial dependence parameter in the L^1 -norm. The number in each subplot represents the proportion of VaR forecast violations




Fig. 2 Rolling Window for $T = 63$ from 2018 to 2020. Rolling window parameter estimation of size $T = 63$ in a data set of size 653 and dimension $n = 50$. 15 different models are considered. The first figure describes the SAR(1) model incorporating only a single general dependence structure (W_G). The last figure depicts the results of the SAR(4) model incorporating the spatial matrices W_G, W_C, W_B, W_I . The number of draws from the limit distribution is set to $B = 300$. The blue line depicts the ratio of the 95%-quantile of the limit distribution given in Lemma (2.6) over the test statistic S from (11). A value below 1 indicates that the assumed model is not correct. The orange line is the accumulated spatial dependence parameter in the L^1 -norm. The number in each subplot represents the proportion of VaR forecast violations

which overlaps with the period of the financial crisis. Thirdly, however, applying the model specification test to the 15 different models also clearly demonstrates that the spatial matrix W_G , which describes a general influence, provides the strongest explanatory power. Country and industry specific effects seem to have a minor impact (W_C and W_B) when considering the ratio and the number of VaR forecast violations (0.1804 for subplot 2 and 0.2372 for subplot 3, respectively). Fourthly, the interaction effect of industry and country affiliation seems to play a minor role in the Euro Stoxx 50 modeling, as both the ratios and the VaR forecast violations of subplots 11 and 15 are similar (0.0546 and 0.0579, respectively). Taking the ratio of VaR forecast violations, the time period in which the null hypothesis cannot be rejected and parsimony in the number of variables as goodness of fit criteria, the SAR(2) incorporating general and industry-specific dependencies, i.e. $\mathbf{y}_t = p_G W_G + p_B W_B + \boldsymbol{\varepsilon}_t$ (cf. subplot 6 of Fig. 1), seems to be the most appropriate.

Figure 2 shows the results of the second application of our proposed model specification test for the period of 2018–2020, where our proceeding is identical to that from the first empirical application: First, we compute the spatial model parameters using a two-step GMM approach. Then, in a second step, we calculate the test statistics of our model specification test. The results of the second application seem to corroborate the findings of the first application: Even if the 15 considered SAR models are rejected by our model specification test in the period 2018–2020, it is striking that models which incorporate at least two spatial matrices model the Euro Stoxx 50 better. This can be seen from the fact that in models with at least two appropriate spatial matrices (W_G should be incorporated) both the ratio of the test statistic and its quantile is larger and VaR forecast violations correspond approximately to the significance level of $\alpha = 5\%$. Furthermore, the beginning of the COVID-19 pandemic in March 2020 is also clearly visible in every subplot of Fig. 2, as the ratio (blue line) decreases significantly. In summary, in terms of parsimony with regard to the number of variables included in a model, the ratio of the test statistic and the quantile and the proportion of VaR forecast violations the SAR(2) model incorporating general and industry-specific effects (cf. subplot 6 in Fig. 2) seems to be the most suitable modeling the Euro Stoxx 50 in the periods considered.

5 Conclusion

In this paper, we propose two novel specification tests for systems of spatial autoregressive models, which are based on the idea of measuring the magnitude of regression residuals. The proposed tests show good size and power properties in finite samples for both initial data and GARCH adjusted data. An empirical analysis of the Euro Stoxx 50 between 2003 to 2009 and 2018 to 2020 gives evidence that a SAR model which models general and industry-specific dependencies is in general appropriate, in particular in bull markets. However, in bear markets a simple spatial model captures the extensive structure of relations and market dependencies to a lesser extent. Accordingly, our proposed testing procedure provides statistical evidence that the model fit is worse in the time after the dot-com bubble, the time around the Lehman Brothers bankruptcy and the time of the COVID-19 pandemic.

In future research, it might be interesting to extend the idea of the tests to other type of spatial dependence models and to compare the fit to stock data with our specific model. Also, a more detailed analysis about the ability of our tests to assess VaR forecast performances or an extension to Δ -hypothesis such as considered in Kutzker et al. (2021) might be interesting.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00181-023-02518-3>.

Funding Open Access funding enabled and organized by Projekt DEAL.

Data availability The stock data used in the empirical application has been downloaded from Thomson Reuters Datastream (first period) and Yahoo! Finance (second period).

Declarations

Conflict of interest The authors have no relevant financial or non-financial interests to declare.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Anselin L (1996) Interactive techniques and exploratory spatial data analysis. Regional Research Institute Working Paper, Nr. 9627
- Arnold M, Stahlberg S, Wied D (2013) Modelling different kinds of spatial dependence in stock returns. *Empir Econ* 44(2):761–774
- Asgharian H, Hess W, Liu L (2013) A spatial analysis of international stock market linkages. *J Bank Finance* 37(12):4738–4754
- Aue A, Hörmann S, Horvath L, Reimherr M (2009) Break detection in the covariance structure of multivariate time series models. *Ann Stat* 37(6B):4046–4087
- Baltagi BH, Song SH, Koh W (2003) Testing panel data regression models with spatial error correlation. *J Econom* 117(1):123–150
- Blasques F, Koopman SJ, Lucas A, Schaumburg J (2016) Spillover dynamics for systemic risk measurement using spatial financial time series models. *J Econom* 195(2):211–223
- Born B, Breitung J (2011) Simple regression based tests for spatial dependence. *Econom J* 14(2):330–342
- Catania L, Billé AG (2017) Dynamic spatial autoregressive models with autoregressive and heteroskedastic disturbances. *J Appl Econom* 32(6):1178–1196
- Demetrescu M, Wied D (2019) Testing for constant correlation of filtered series under structural change. *Econom J* 22(1):10–33
- Fernández-Avilés G, Montero JM, Orlov AG (2012) Spatial modeling of stock market comovements. *Finance Res Lett* 9(4):202–212
- Fleming MM (2004) Techniques for estimating spatially dependent discrete choice models. In: Anselin L, Florax R, Rey SJ (eds) *Advances in spatial econometrics: methodology, tools and applications*. Springer, Berlin
- Fülle MJ, Otto P (2023) Spatial GARCH models for unknown spatial locations—an application to international stock returns. *Spatial Econ Anal* (forthcoming)

- Gong P, Weng Y (2016) Value-at-risk forecasts by a spatiotemporal model in Chinese stock market. *Physica A Stat Mech Appl* 441(C):173–191
- Hansen LP (1982) Large sample properties of generalized method of moments estimators. *Econometrica* 50:1029–1054
- Herrera M, Mur J, Ruiz M (2019) A comparison study on criteria to select the most adequate weighting matrix. *Entropy* 21(2):160
- Kelejian HH, Piras G (2011) An extension of Kelejian's J-test for non-nested spatial models. *Reg Sci Urban Econ* 41(3):281–292
- Kelejian HH, Piras G (2016) An extension of the J-test to a spatial panel data framework. *J Appl Econom* 31(2):387–402
- Kelejian HH, Prucha IR (2001) On the asymptotic distribution of the Moran I test statistic with applications. *J Econom* 104(2):219–257
- Kutzker T, Stark F, Wied D (2021) Testing for relevant dependence change in financial data: a CUSUM copula approach. *Emp Econ* 60(4):1875–1894
- LeSage JP, Pace RK (2014) The biggest myth in spatial econometrics. *Econometrics* 2(4):217–249
- Li H, Calder CA, Cressie N (2007) Beyond Moran's I: Testing for spatial dependence based on the spatial autoregressive model. *Geogr Anal* 39(4):357–375
- Mattera R, Otto P (2023) Network log-arch models for forecasting stock market volatility. [arXiv:2303.11064](https://arxiv.org/abs/2303.11064)
- Millo G (2017) A simple randomization test for spatial correlation in the presence of common factors and serial correlation. *Reg Sci Urban Econ* 66:28–38
- Moran PAP (1950) Notes on continuous stochastic phenomena. *Biometrika* 37(1/2):17–23
- Piras G, Elhorst JP, Lacombe DJ (2012) On model specification and parameter space definitions in higher order spatial econometric models. *Reg Sci Urban Econ* 42:211–220
- Schmitt T, Schäfer R, Wied D, Guhr T (2016) Spatial dependence in stock returns—local normalization and VaR forecasts. *Empir Econ* 50(3):1091–1109
- Selan B, Kalatzis AEG (2017) Peer effects of stock returns and financial characteristics: spatial approach for an emerging market. Working Paper
- Su L, Qu X (2017) Specification tests for spatial autoregressive models. *J Bus Econ Stat* 35(4):572–584
- Tam PS (2014) A spatial-temporal analysis of east Asian equity market linkages. *J Comp Econ* 42(2):304–327
- Wied D (2013) CUSUM-type testing for changing parameters in a spatial autoregressive model for stock returns. *J Time Ser Anal* 34(1):211–229
- Zhang W-G, Mo G-L, Liu F, Liu Y-J (2018) Value-at-risk forecasts by dynamic spatial panel GJR-GARCH model for international stock indices portfolio. *Soft Comput* 22:5279–5297
- Zhou J, Li D, Pan R, Wang H (2020) Network GARCH model. *Stat Sin* 30(4):1723–1740
- Ziggel D, Berens T, Weiß G, Wied D (2014) A new set of improved value-at-risk backtests. *J Bank Finance* 48:29–41